

A FAMILY OF MULTIPLE HARMONIC SUM AND MULTIPLE ZETA STAR VALUE IDENTITIES

ERIN LINEBARGER & JIANQIANG ZHAO

ABSTRACT. In this paper we present a new family of identities for multiple harmonic sums which generalize a recent result of Hessami Pilehrood et al. [1]. We then apply it to obtain a family of identities of multiple zeta star values.

1. INTRODUCTION

In a recent paper [1] Hessami Pilehrood et al. proved a few families of identities involving (alternating) multiple harmonic sums and binomial coefficients and, as applications, discovered some new congruences for multiple harmonic sums. In particular, they are able to confirm several conjectures contained in [4] posed by the second author of this paper. They can also provide some new families of identities of MZSV and a new proof of the identity of Zagier [3].

To state one of their results we recall that the (alternating) multiple harmonic sums are multiple variable generalizations of harmonic sums defined by the following: for any $r \in \mathbb{N}$ and $\mathbf{s} = (s_1, s_2, \dots, s_r) \in (\mathbb{Z}^*)^r$

$$H_n(s_1, s_2, \dots, s_r) = \sum_{n \geq k_1 > k_2 > \dots > k_r \geq 1} \prod_{i=1}^r \frac{\text{sgn}(s_i)^{k_i}}{k_i^{|s_i|}}.$$

Another kind of sums which is closely related to the above is the following star version (also denoted by S in the literature)

$$H_n^*(s_1, s_2, \dots, s_r) = \sum_{n \geq k_1 > k_2 > \dots > k_r \geq 1} \prod_{i=1}^r \frac{\text{sgn}(s_i)^{k_i}}{k_i^{|s_i|}}.$$

We call $l(\mathbf{s}) := r$ the depth of this MHS and $|\mathbf{s}| := \sum_{i=1}^r |s_i|$ the weight. For convenience we set $H_n(\mathbf{s}) = 0$ if $n < l(\mathbf{s})$, $H_n(\emptyset) = H_n^*(\emptyset) = 1$, and $\{s_1, s_2, \dots, s_j\}^d$ the set formed by repeating the composition $(s_1, s_2, \dots, s_j)$ d times. When taking the limit $n \rightarrow \infty$ we get the so-called the (alternating) Euler sum and the (alternating) star Euler sum, respectively:

$$\zeta(\mathbf{s}) = \lim_{n \rightarrow \infty} H_n(\mathbf{s}), \quad \zeta^*(\mathbf{s}) = \lim_{n \rightarrow \infty} H_n^*(\mathbf{s}). \quad (1)$$

1991 *Mathematics Subject Classification.* 11A07, 11M32, 11B65, 11B68.

Key words and phrases. Multiple harmonic sum, multiple zeta star values.

When $\mathbf{s} \in \mathbb{N}^\ell$ they are called the *multiple zeta value* (MZV) and the *multiple zeta star value* (MZSV), respectively.

The following is one of the main results of [1].

Theorem 1.1. ([1, Thm. 2.1]) *Let a and b be two non-negative integers. Set $A_{n,k} = (-1)^{k-1} \binom{n}{k} / \binom{n+k}{k}$ for any positive integers n and k . Then for any integer $c \geq 2$,*

$$H_n^*(\{2\}^a, c, \{2\}^b) = 2 \sum_{k=1}^n \frac{A_{n,k}}{k^{2a+2b+c}} + 4 \sum_{\substack{i+j+|\mathbf{x}|=c \\ i \geq 1, j \geq 2}} 2^{l(\mathbf{x})} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{x}, i+2b) A_{n,k}}{k^{2a+j}}. \quad (2)$$

In sections 2 and 3 we generalize Theorem 1.1 to express $H_n^*(\mathbf{s})$ using binomial coefficients where $\mathbf{s} = (\{2\}^{a_1}, c_1, \dots, \{2\}^{a_r}, c_r, \{2\}^{a_{r+1}})$ with positive integers $c_1, \dots, c_r \geq 2$. Taking limit and using a key lemma we shall obtain a new family of identities of MZSV involving alternating Euler sums in section 4.

The ideas used in this paper have been applied to obtain similar results for other types of strings in [5]. In particular, the conjectural Two-one formula of Ohno and Zudlin is proved there. In [2] Hessami Pilehroods and the second author will further prove a q -analog of the main result of this paper (Theorem 2.3) and some similar results for other types of strings contained in [5].

2. IDENTITIES FOR MULTIPLE HARMONIC SUMS

The following lemma is the special case of [1, Lemma 2.2] when $m = 2$ and $c_n^{(2)} = n!^2/(2n)!$.

Lemma 2.1. *For any positive integers n and k define $A_{n,k} = (-1)^{k-1} \binom{n}{k} / \binom{n+k}{k}$. Then for every $c \in \mathbb{N}$ and $\mathbf{v} \in \mathbb{N}^t$ ($t \geq 0$) we have*

$$\frac{1}{n^c} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{v}) A_{n,k}}{k^a} = \sum_{k=1}^n \frac{H_{k-1}(\mathbf{v}) A_{n,k}}{k^{a+c}} + \sum_{\substack{j+|\mathbf{x}|=a+c \\ j \geq 0, x_r > a}} 2^{l(\mathbf{x})} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{x}, \mathbf{v}) A_{n,k}}{k^j} \quad (3)$$

where $\mathbf{x} = (x_1, x_2, \dots, x_r) \in \mathbb{N}^r$ for $r \geq 0$, and $|\mathbf{x}| = \sum_{i=1}^r x_i$, $l(\mathbf{x}) = r$. Here $|\mathbf{x}| = 0$ if $r = 0$ which implies that $\mathbf{x} = \emptyset$.

Definition 2.2. Suppose $\mathbf{s} = (\{2\}^{a_1}, c_1, \dots, \{2\}^{a_r}, c_r, \{2\}^{a_{r+1}})$. We define two kinds of string operations on the *condensation* $\tilde{\mathbf{s}} := (2a_1, c_1, \dots, 2a_r, c_r, 2a_{r+1})$ of $\mathbf{s}$ as follows.

- A *merge* μ_t for some $1 \leq t \leq r$ changes the symbols “ c_t ,” to “ $+c_t+$ ”,
- A *substitution* σ_t changes the symbols “ c_t ,” to “ $+j_t, \mathbf{x}_t, i_t+$ ”.

Let $I = \{i_1, \dots, i_t\}$ be a subset of $[r] := \{1, \dots, r\}$. Define

- The merge operation $\mu_I = \mu_{i_1} \circ \dots \circ \mu_{i_t}$ and similarly for substitutions σ_I ,
- The complement $\bar{I} = [r] \setminus I$,

- The combined operation $\kappa_I = \sigma_I \circ \mu_I$.

For any composition $(e_1, \dots, e_r)$ of natural numbers we denote

- The first component $e_1 = \varphi(\mathbf{e})$
- The truncation operator $\text{Tr}(e_1, \dots, e_r) = (e_2, \dots, e_r)$.
- The negation operator $\nu(e_1, \dots, e_r) = (\bar{e}_1, e_2, \dots, e_r)$.

Notice that if $r = 1$ then $\text{Tr}(e_1) = \emptyset$.

Theorem 2.3. *Let n be a positive integer and $\mathbf{s} = (\{2\}^{a_1}, c_1, \dots, \{2\}^{a_r}, c_r, \{2\}^{a_{r+1}})$ where $a_j, c_j \in \mathbb{N}_0$ and $c_j \geq 2$. Set $A_{n,k} = (-1)^{k-1} \binom{n}{k} / \binom{n+k}{k}$. Then*

$$H_n^*(\mathbf{s}) = 2 \sum_{I \subseteq [r]} \sum_{\substack{i_t + j_t + |\mathbf{x}_t| = c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1}(\text{Tr} \circ \kappa_I(\tilde{\mathbf{s}})) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{s}})}}, \quad (4)$$

where $\tilde{\mathbf{s}} = (2a_1, c_1, \dots, 2a_r, c_r, 2a_{r+1})$ is the condensation of $\mathbf{s}$.

Remark 2.4. When $r = 1$ we have two possible subsets of $[1]$: $I = \emptyset$ and $I = [1]$. our Theorem 2.3 becomes Theorem 1.1. When $r = 2$ we get the following: For a positive integer n and non-negative integers a_1, a_2, a_3, c_1, c_2 we have

$$\begin{aligned} & H_n^*(\{2\}^{a_1}, c_1, \{2\}^{a_2}, c_2, \{2\}^{a_3}) \\ &= 2 \sum_{k=1}^n \frac{A_{n,k}}{k^{2a_1+c_1+2a_2+c_2+2a_3}} + 4 \sum_{\substack{i+j+|\mathbf{x}|=c_2 \\ i \geq 1, j \geq 2}} 2^{l(\mathbf{x})} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{x}, i+2a_3) A_{n,k}}{k^{2a_1+c_1+2a_2+j}} \\ &+ 8 \sum_{\substack{i_1+j_1+|\mathbf{x}_1|=c_1, i_2+j_2+|\mathbf{x}_2|=c_2 \\ i_1, i_2 \geq 1, j_1, j_2 \geq 2}} 2^{l(\mathbf{x}_1)+l(\mathbf{x}_2)} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{x}_1, i_1+2a_2+j_2, \mathbf{x}_2, i_2+2a_3) A_{n,k}}{k^{2a_1+j_1}} \\ &+ 4 \sum_{\substack{i+j+|\mathbf{x}|=c_1 \\ i \geq 1, j \geq 2}} 2^{l(\mathbf{x})} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{x}, i+2a_2+c_2+2a_3) A_{n,k}}{k^{2a_1+j}}. \end{aligned} \quad (5)$$

3. PROOF OF THEOREM 2.3

We proceed by induction on the sum $r + n$. Since $r \geq 0, n \geq 1$, our base case is $r + n = 1$ which implies $H_1^*(\{2\}^a) = 1$, and therefore the formula is true. Suppose the statement is true when $r + n = k - 1$, and let $r + n = k$. Then by definition

$$H_n^*(\mathbf{s}) = \sum_{l=0}^{a_1} \frac{1}{n^{2(a_1-l)}} H_{n-1}^*(\mathbf{t}_l) + \frac{1}{n^{2a_1+c_1}} H_n^*(\mathbf{u}),$$

where $\mathbf{t}_l = (\{2\}^l, c_1, \{2\}^{a_2}, c_2, \dots, c_r, \{2\}^{a_{r+1}})$ and $\mathbf{u} = (\{2\}^{a_2}, c_2, \dots, c_r, \{2\}^{a_{r+1}})$. We now set their condensations as follows:

$$\tilde{\mathbf{t}}_l = (2l, c_1, 2a_2, c_2, \dots, c_r, 2a_{r+1}), \quad \tilde{\mathbf{u}} = (2a_2, c_2, \dots, c_r, 2a_{r+1}).$$

By inductive hypothesis,

$$\begin{aligned} H_n^*(\mathbf{s}) &= \sum_{l=0}^{a_1} \frac{2}{n^{2(a_1-l)}} \sum_{I \subseteq [r]} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^{n-1} \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l) \right) A_{n-1,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l)}} \\ &\quad + \frac{2}{n^{2a_1+c_1}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{u}}) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{u}})}}. \end{aligned}$$

Changing the order of summation and summing the inner sum we get

$$A_{n-1,k} \sum_{l=0}^{a_1} \frac{n^{2l}}{k^{2l}} = (-1)^{k-1} \frac{n^{2a_1+2} - k^{2a_1+2}}{(n^2 - k^2)k^{2a_1}} \frac{\binom{n-1}{k}}{\binom{n-1+k}{k}} = \left(\frac{n^{2a_1}}{k^{2a_1}} - \frac{k^2}{n^2} \right) A_{n,k}.$$

Thus

$$\begin{aligned} H_n^*(\mathbf{s}) &= 2 \sum_{I \subseteq [r]} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l) - 2l + 2a_1}} \\ &\quad - \frac{2}{n^{2a_1+2}} \sum_{I \subseteq [r]} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l) - 2l - 2}} \\ &\quad + \frac{2}{n^{2a_1+c_1}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{u}}) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{u}})}}. \end{aligned}$$

Since $\tilde{\mathbf{t}}_l$ coincides with $\tilde{\mathbf{s}}$ everywhere except for the first component, $\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l) = \text{Tr} \circ \kappa_I(\tilde{\mathbf{u}})$. Similarly, $\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l) - 2l + 2a_1 = \varphi \circ \kappa_I(\tilde{\mathbf{u}})$. Hence, by induction we only need to show the following quantity vanishes:

$$\begin{aligned} &\frac{2}{n^{2a_1+c_1}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{u}}) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{u}})}} \\ &- \frac{2}{n^{2a_1+2}} \sum_{I \subseteq [r]} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l) - 2l - 2}} \end{aligned}$$

$$\begin{aligned}
&= \frac{2}{n^{2a_1+c_1}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{u}}) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{u}})}} \\
&\quad - \frac{2}{n^{2a_1+2}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l) - 2l - 2}} \\
&\quad - \frac{4}{n^{2a_1+2}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_{I \cup \{1\}}(\tilde{\mathbf{t}}_l) \right) A_{n,k}}{k^{\varphi \circ \mu_{\bar{I}} \circ \sigma_{I \cup \{1\}}(\tilde{\mathbf{t}}_l) - 2l - 2}} \\
&= \frac{2}{n^{2a_1+2}} \sum_{I \subseteq [r] \setminus \{1\}} \sum_{\substack{i_t+j_t+|\mathbf{x}_t|=c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \left(\frac{1}{n^{c_1-2}} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{u}}) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{u}})}} \right. \\
&\quad \left. - 2 \sum_{\substack{i_1+j_1+|\mathbf{x}_1|=c_1, \\ i_1 \geq 1, j_1 \geq 2}} 2^{l(\mathbf{x}_1)} \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_I(\tilde{\mathbf{t}}_l) \right) A_{n,k}}{k^{\varphi \circ \kappa_I(\tilde{\mathbf{t}}_l) - 2l - 2}} - \sum_{k=1}^n \frac{H_{k-1} \left(\text{Tr} \circ \kappa_{I \cup \{1\}}(\tilde{\mathbf{t}}_l) \right) A_{n,k}}{k^{\varphi \circ \kappa_{I \cup \{1\}}(\tilde{\mathbf{t}}_l) - 2l - 2}} \right). \tag{6}
\end{aligned}$$

Note that if $1 \notin I$ then $\text{Tr} \circ \kappa_I(\tilde{\mathbf{u}}) = \text{Tr} \circ \kappa_{I \cup \{1\}}(\tilde{\mathbf{t}}_l)$, which we will now denote by $\mathbf{v}$. Furthermore, $\varphi \circ \kappa_{I \cup \{1\}}(\tilde{\mathbf{t}}_l) = \varphi \circ \kappa_I(\tilde{\mathbf{u}}) + c_1 + 2l$. Let $a = \varphi \circ \kappa_I(\tilde{\mathbf{u}})$. Then

$$\kappa_I(\tilde{\mathbf{t}}_l) = \sigma_{\bar{I}} \circ \mu_I(\tilde{\mathbf{t}}_l) = \sigma_1(2l, c_1, a, \dots) = (j_1 + 2l, \mathbf{x}_1, i_1 + a, \dots).$$

Set $\mathbf{w} = (\mathbf{x}_1, i_1 + a)$. Then we see that the three sums inside the parenthesis (6) can be rewritten as

$$\frac{1}{n^{c_1-2}} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{v}) A_{n,k}}{k^a} - \sum_{\substack{j_1+|\mathbf{w}|=c_1+a_1, \\ j_1 \geq 2, w_p > a_1}} 2^{l(\mathbf{w})} \sum_{k=1}^n \frac{H_{k-1}(\mathbf{w}, \mathbf{v}) A_{n,k}}{k^{j_1-2}} - \sum_{k=1}^n \frac{H_{k-1}(\mathbf{v}) A_{n,k}}{k^{a+c_1-2}},$$

where $w_p = i + a_1$ ($i \geq 1$) is the last component of $\mathbf{w}$. This last expression vanishes by taking $c = c_1 - 2$ and $j = j_1 - 2$ in Lemma 2.1. We have now completed the proof of Theorem 2.3.

4. THE KEY LEMMA AND AN IDENTITY FAMILY OF MZSV

In [1], using the corresponding identities of MHS Hessami Pilehrood et al. find some new families of identities of MZSV and a new proof of the identity of Zagier [3]. We can similarly use Theorem 2.3 to obtain a new family of MZSV as follows.

Theorem 4.1. *Let n be a positive integer and $\mathbf{s} = (\{2\}^{a_1}, c_1, \dots, \{2\}^{a_r}, c_r, \{2\}^{a_{r+1}})$ where $a_j, c_j \in \mathbb{N}_0$ and $c_j \geq 2$. Set $\tilde{\mathbf{s}} = (2a_1, c_1, \dots, 2a_r, c_r, 2a_{r+1})$. Then with notation given by Definition 2.2 we have*

$$\zeta^*(\mathbf{s}) = 2 \sum_{I \subseteq [r]} \sum_{\substack{i_t + j_t + |\mathbf{x}_t| = c_t, \\ i_t \geq 1, j_t \geq 2 \quad \forall t \notin I}} 2^{|\bar{I}| + \sum_{t \notin I} l(\mathbf{x}_t)} \zeta\left(\nu \circ \kappa_I(\tilde{\mathbf{s}})\right). \quad (7)$$

Proof. By taking limit $n \rightarrow \infty$ in Theorem 2.3 we see that (7) follows immediately from the following lemma. $\square$

Lemma 4.2. *Let $d \in \mathbb{N}_0$ and let e be a real number with $e > 1$. Then for all $\mathbf{s} \in (\mathbb{Z}^*)^d$ ($\mathbf{s} = \emptyset$ if $d = 0$) we have*

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{|H_{k-1}(\mathbf{s})|}{k^e} \left(1 - \frac{\binom{n}{k}}{\binom{n+k}{k}}\right) = 0. \quad (8)$$

Remark 4.3. This is the key step which enables us to go from MHS identities to MZSV identities in this paper. Apparently the authors of [1] have already used this result in their paper although no proof is given there. Because of its importance we provide the following detailed analysis. Notice that for any fixed k we have $\lim_{n \rightarrow \infty} \binom{n}{k} / \binom{n+k}{k} = 1$ but for k close to n , say $k = n$, we have $\lim_{n \rightarrow \infty} \binom{n}{k} / \binom{n+k}{k} = 0$ by Sterling's formula. So (8) is not obvious to us.

Proof. First we have

$$\begin{aligned} 0 < 1 - \frac{\binom{n}{k}}{\binom{n+k}{k}} &= 1 - \frac{n(n-1)(n-2) \cdots (n-k+1)}{(n+1)(n+2) \cdots (n+k)} \\ &= 1 - \frac{n(n-1)(\frac{n}{2}-1) \cdots (\frac{n}{k-1}-1)}{k(n+1)(\frac{n}{2}+1) \cdots (\frac{n}{k}+1)} \\ &= 1 - \frac{n^k H_{k-1}(\{1\}^{k-1}) - n^{k-1} H_{k-1}(\{1\}^{k-2}) + \cdots + (-1)^k n^2 H_{k-1}(1) - n(-1)^k}{k(n^k H_k(\{1\}^k) + n^{k-1} H_k(\{1\}^{k-1}) + \cdots + n H_k(1) + 1)} \end{aligned} \quad (9)$$

$$\begin{aligned} &\leq \frac{n^{k-1} H_k(\{1\}^{k-2}) + n^{k-2} H_k(\{1\}^{k-3}) + \cdots + n^2 H_k(1) + n}{k(n^k H_k(\{1\}^k) + n^{k-1} H_k(\{1\}^{k-1}) + \cdots + n H_k(1) + 1)} \\ &\quad + \frac{n^{k-1} H_k(\{1\}^{k-1}) + \cdots + n H_k(1) + 1}{n^k H_k(\{1\}^k) + n^{k-1} H_k(\{1\}^{k-1}) + \cdots + n H_k(1) + 1}. \end{aligned} \quad (10)$$

Here from (9) to (10) we canceled the leading term of the numerator in (9) because of $H_k(\{1\}^k) = 1/k!$, took absolute values of each term on the right and then changed all

H_{k-1} to H_k . Observe that for any $1 \leq j \leq k$ we have

$$\begin{aligned} H_k(\{1\}^j) &= \sum_{k \geq n_1 > \dots > n_j > 0} \frac{1}{n_1 \cdots n_j} \\ &\leq \sum_{k \geq n_1 > \dots > n_j > n_{j+1} > 0} \frac{k}{n_1 \cdots n_l \cdot n_{j+1}} = k H_k(\{1\}^{j+1}) \leq \dots \leq k^{k-j} H_k(\{1\}^k), \end{aligned}$$

which is even true for $j = 0$. Thus we get

$$1 - \frac{\binom{n}{k}}{\binom{n+k}{k}} \leq \frac{2(n^{k-1}k + n^{k-2}k^2 + \dots + nk^{k-1} + k^k)}{n^k} \leq \frac{2k^2}{n}.$$

Now for any $\mathbf{s} \in (\mathbb{Z}^*)^d$ we clearly have

$$|H_{k-1}(\mathbf{s})| \leq H_{k-1}(\{1\}^d) \leq C \log^d(k)$$

for some positive constant C . Therefore by setting $\delta = \min\{(e-1)/2, 1/2\} > 0$ we have

$$\sum_{k=1}^n \frac{|H_{k-1}(\mathbf{s})|}{k^e} \left(1 - \frac{\binom{n}{k}}{\binom{n+k}{k}}\right) \leq \frac{2C}{n} \sum_{k=1}^{\lfloor n^\delta \rfloor} \frac{\log^d(k)}{k^{e-2}} + C \sum_{k=\lfloor n^\delta \rfloor+1}^{\infty} \frac{\log^d(k)}{k^e}. \quad (11)$$

If $e \geq 2$ then

$$\frac{2C}{n} \sum_{k=1}^{\lfloor n^\delta \rfloor} \frac{\log^d(k)}{k^{e-2}} \leq \frac{2C}{n} \sum_{k=1}^{\sqrt{n}} \log^d(k) \leq \frac{2C \log^d(\sqrt{n})}{\sqrt{n}} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

If $1 < e < 2$ then

$$\frac{2C}{n} \sum_{k=1}^{\lfloor n^\delta \rfloor} \frac{\log^d(k)}{k^{e-2}} \leq \frac{2C n^{\delta(3-e)} \log^d(n^\delta)}{n} \leq \frac{2C \delta^d \log^d(n)}{\sqrt{n}} \rightarrow 0 \text{ as } n \rightarrow \infty,$$

since $1 + \delta(e-3) = 1 + (e-1)(e-3)/2 = (e-2)^2/2 + 1/2 \geq 1/2$. Now the last sum on the right of (11) is the tail of a convergent series so (8) follows immediately and therefore the lemma is proved. $\square$

Acknowledgement. We would like to thank Roberto Tauraso for sending us their preprint [1] upon which we can build our current work, Wadim Zudilin for kindly showing us an improvement on Lemma 4.2, and K. Hessami Pilehrood and T. Hessami Pilehrood for pointing out some misprints in the first version of this paper. JZ is supported by an NSF grant DMS1162116.

REFERENCES

- [1] K. Hessami Pilehrood, T. Hessami Pilehrood and R. Tauraso, New properties of multiple harmonic sums modulo p and p -analogues of Leshchiner's series, to appear in *Trans. Amer. Math. Soc.*, arXiv:1206.0407
- [2] K. Hessami Pilehrood, T. Hessami Pilehrood and J. Zhao, On q -analogues of some identity families for multiple harmonic sums and multiple zeta star values, in preparation.
- [3] D. Zagier, Evaluation of the multiple zeta values $\zeta(2, \dots, 2, 3, 2, \dots, 2)$, *Ann. of Math.* **175** (2012), pp. 977–1000.
- [4] J. Zhao, Wolstenholme type Theorem for multiple harmonic sums, *Int. J. of Number Theory* **4** (2008), no. 1, pp. 73–106, arXiv:math.NT/0301252
- [5] J. Zhao, Identity families of multiple harmonic sums and multiple zeta (star) values, preprint arXiv:1303.2227

DEPARTMENT OF MATHEMATICS, ECKERD COLLEGE, ST. PETERSBURG, FL 33711

E-mail address: emlineba@eckerd.edu

E-mail address: zhaoj@eckerd.edu